

THE STELLAR MASSES OF DISK GALAXIES AND THE CALIBRATION OF COLOR-MASS-TO-LIGHT RATIO RELATIONS

STACY S. MCGAUGH

Department of Astronomy, Case Western Reserve University, Cleveland, OH 44106

AND

JAMES M. SCHOMBERT

Department of Physics, University of Oregon, Eugene, OR 97403

Draft version March 5, 2013

ABSTRACT

We present new Spitzer 3.6 μ m observations of a sample of disk galaxies spanning over 10 magnitudes in luminosity and ranging in gas fraction from $\sim 10\%$ to over 90%. We use these data to test population synthesis prescriptions for computing stellar mass. Many commonly employed models fail to provide self-consistent stellar masses in the sense that the stellar mass estimated from the optical luminosity typically exceeds that estimated from the near-infrared (NIR) luminosity. This problem is present in models both with and without TP-AGB stars, but is more severe in the former. Self-consistency can be achieved if NIR mass-to-light ratios are approximately constant with a mean value near $0.5 M_{\odot}/L_{\odot}$ at 3.6 μ m. We use the Baryonic Tully-Fisher relation calibrated by gas rich galaxies to provide an independent estimate of the color-mass-to-light ratio relation. This approach also suggests that the typical 3.6 μ m mass-to-light ratio is $0.5 M_{\odot}/L_{\odot}$ ($0.65 M_{\odot}/L_{\odot}$ in the K_s -band) for rotationally supported galaxies. These values are consistent with a Kroupa IMF.

Subject headings: galaxies: evolution — galaxies: fundamental parameters — galaxies: photometry
 — galaxies: kinematics and dynamics — galaxies: stellar content

1. INTRODUCTION

One of the most fundamental properties of a galaxy is its luminosity and the mass of the stars that produce it. Our understanding of stellar evolution is sufficiently advanced that it should be possible to compute the luminosity produced by a stellar population *ab initio*. Indeed, there exist in the literature various prescriptions for estimating stellar mass from observed colors or spectral energy distributions (e.g., Bell & de Jong 2001; Bell et al. 2003; Portinari et al. 2004; Zibetti et al. 2009; Into & Portinari 2013).

A general expectation of population synthesis models is that the relation between mass and light is more nearly constant in the near-infrared (NIR) than in the optical part of the spectrum. This follows from basic considerations: recent star formation populates the upper main sequence with luminous, blue stars. These stars produce copious amounts of optical light from little mass and lead short lives, leading to substantial perturbations on the average mass-to-light ratio Υ_* of a galaxy. The degree to which this occurs depends on the IMF and the intensity of a star forming event relative to the total stellar mass already present. These effects combine to make the prediction of any particular galaxy's optical mass-to-light ratio uncertain at the factor of a few level. Young stars contribute rather less in the NIR part of the spectrum, so one expects a closer relation between light and mass at these wavelengths. Empirically, the scatter in the Tully-Fisher relation declines as one goes from blue to red to NIR wavelengths (Verheijen 2001), consistent with the expected decrease in scatter in Υ_* .

In this paper we present new 3.6 μ m Spitzer Space Telescope photometry of a sample of galaxies spanning a large range (ten magnitudes) in luminosity. This wavelength is expected to be close to the sweet spot of nearly constant Υ_* . We combine these data with optical colors and luminosities to check the predictions of several population synthesis models.

We describe the data in §2. In §3 we compute stellar masses with various population synthesis models, and estimate the mean NIR mass-to-light ratio that is required for each model to be self-consistent. In §4 we utilize the Baryonic Tully-Fisher Relation (BTFR) to obtain an independent, kinematic estimate of stellar mass. We discuss our results in §5, including some of the implications for galaxy gas fractions in §5.1. We summarize our conclusions in §6.

2. DATA

The SINGS (Kennicutt et al. 2003; Dale et al. 2005) and THINGS (Walter et al. 2008; de Blok et al. 2008) surveys have demonstrated the utility of Spitzer 3.6 μ m data for constraining the stellar components of star forming galaxies. Here we wish to sample disk galaxies over as large a range of physical properties as possible. To this end, we combine the THINGS data of de Blok et al. (2008) with new data from from two Spitzer programs. One cycle 5 project targeted galaxies with high quality rotation curves selected to increase the sampling of circular velocities to both higher and lower mass than already present in THINGS. A cycle 7 snapshot program provides additional photometry for low surface brightness galaxies. The combined sample spans ten magnitudes in [3.6] luminosity. All galaxies have at least some ancillary optical photometry, and a large subset also has kinematic information that can be used for the Tully-Fisher rela-

Table 1
Galaxy Data

Galaxy	D	M_V	$M_{[3.6]}$	$B - V$	$V - I$	$J - K$	Refs.
D564-8	6.51	-12.64	-15.14	...	0.93	...	1,3
KK98 251	6.08	...	-16.39	...	...	...	1
DDO 154	4.04	-14.55	-16.41	0.32	...	0.34	2,3
D631-7	5.49	-14.50	-16.73	0.41	0.55	...	1,4
D640-13	13.44	-14.36	-16.77	...	0.55	...	1,4
D568-2	21.3	-14.6	-16.81	0.45	0.70	...	1,3
D572-5	14.56	-14.56	-16.82	0.44	0.52	...	1,4
F415-3	10.4	-15.2	-16.99	0.62	...	...	1,5
D500-3	18.52	-15.74	-17.18	...	0.31	...	1,4
DDO 168	4.25	-15.70	-17.45	0.32	...	...	1,3
D575-1	9.79	-14.22	-17.51	...	0.70	...	1,4
D512-2	14.12	-15.22	-17.83	...	0.80	...	1,4
F611-1	25.5	-15.4	-17.92	0.57	...	...	1,5
D500-2	17.89	-16.38	-18.25	0.52	0.42	...	1,4
DDO 170	16.3	...	-18.50	...	...	...	1
F565-V2	55.1	-16.2	-18.76	0.44	...	...	1,5
NGC 2366	3.27	-17.71	-18.90	0.54	...	0.84	2
D723-5	27.7	-16.9	-19.21	0.55	0.75	...	1,3
F563-V1	57.6	-17.2	-19.79	0.23	0.83	...	1,5
IC 2574	3.91	-18.53	-20.15	0.42	...	0.58	2
F563-1	52.2	-17.8	-20.40	0.40	0.86	...	1,5
F574-2	92.3	-18.3	-20.50	0.58	...	...	1,5
NGC 2976	3.58	-18.33	-20.52	0.55	...	0.79	2
F568-V1	84.8	-18.6	-20.82	0.47	...	...	1,5
F561-1	69.8	-18.3	-20.88	0.69	...	...	1,5
F577-V1	113.	-18.7	-20.95	0.50	...	...	1,5
NGC 1003	10.24	-20.38	-21.12	0.42	...	0.73	2
UGC 5005	57.1	-18.7	-21.18	0.35	...	...	1,5
F574-1	100.	-19.1	-21.37	0.51	...	...	1,5
F568-1	95.5	-18.9	-21.38	0.52	...	...	1,5
NGC 7793	3.61	-19.42	-21.46	0.63	...	0.68	2
UGC 128	58.5	-19.3	-21.88	0.63	0.68	...	1,5
NGC 2403	3.16	-19.82	-21.97	0.39	...	0.75	2
NGC 925	9.43	-20.54	-22.30	0.50	...	0.83	2
NGC 2903	8.9	-21.48	-22.74	0.55	...	0.91	2
NGC 3198	13.8	-21.18	-23.00	0.43	...	0.92	2
NGC 3621	6.56	-20.57	-23.04	0.52	...	0.83	2
NGC 3521	8.0	-21.62	-24.19	0.68	...	0.93	2
NGC 3031	3.65	-21.55	-24.28	0.82	...	0.88	2
NGC 5055	8.99	-21.76	-24.60	0.64	...	0.95	2
NGC 2998	68.3	-22.36	-24.77	0.45	...	0.99	2
NGC 2841	14.1	-21.95	-24.88	0.74	...	0.93	2
UGC 2953	15.4	...	-25.08	...	...	0.90	1
NGC 6674	51.9	-22.17	-25.17	0.57	...	0.86	2
NGC 7331	14.9	-22.30	-25.30	0.63	...	1.00	2
NGC 801	75.3	-22.30	-25.33	0.61	...	1.05	2
NGC 5533	59.4	-22.16	-25.47	0.77	...	0.94	2
UGC11455	75.4	...	-25.61	...	...	1.16	1
UGCA 145	64.5	...	-25.61	...	...	1.10	1
UGC 2885	75.9	-23.30	-25.94	0.47	...	0.88	1

References. — Spitzer [3.6] magnitudes: 1. This work 2. de Blok et al. (2008). Ancillary data: 3. Schombert et al. (2011) 4. Trachternach et al. (2009) 5. McGaugh & de Blok (1998).

Note. — Galaxy photometric data in order of increasing [3.6] luminosity. Adopted distances are in Mpc. $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is assumed when no direct determination is available. Absolute magnitudes assume $M_{\odot}^V = 4.83$ and $M_{\odot}^{[3.6]} = 3.24$.

tion.

The [3.6] luminosities of galaxies in the THINGS sample have been adopted from the mass models of de Blok et al. (2008). Only the total luminosity is used here; no distinction is made between bulge and disk components. For the remainder of the assembled sample, the Spitzer data have been analyzed with the ARCHANGEL surface photometry package (Schombert 2011). Elliptical isophotes have been fit to the data and integrated magnitudes determined from asymptotic fits to curves of growth. Special care has been taken to exclude fore-

ground stars and replace the masked region with an estimate of the galaxy light based on surrounding pixels. IRAC is a sensitive instrument, and many foreground stars (presumably copious numbers of M dwarfs) are detected that are not obvious in survey material. Careful cleaning of these contaminants is essential to accurate photometry.

The data are presented in Table 1. For each galaxy, we assign a distance and compute the corresponding absolute magnitude in the optical V -band and NIR Spitzer IRAC [3.6] band. Distances are taken from the direct

Table 2
Population Synthesis Mass-to-Light Ratios

Model	IMF	a_V	b_V	a_K	b_K	$\Upsilon_{0.6}^V$	$\Upsilon_{0.6}^K$	$\Upsilon_{0.6}^{[3.6]}$	TP
Bell & de Jong (2001)	Scaled Salpeter	−0.734	1.404	−0.692	0.652	1.28	0.50	0.41	no
Bell et al. (2003)	Scaled Salpeter	−0.628	1.305	−0.206	0.135	1.43	0.73	0.62	no
Portinari et al. (2004)	Kroupa (2002)	−0.654	1.290	−0.736	0.730	1.32	0.50	0.41	no
Zibetti et al. (2009)	Chabrier (2003)	−1.075	1.837	−1.390	1.176	1.07	0.21	0.14	yes
Into & Portinari (2013)	Kroupa (2002)	−0.900	1.627	−1.020	1.054	1.19	0.41	0.33	yes

Note. — Stellar mass-to-light ratios in the V and K -bands from various population synthesis models are given in solar units by the formula $\log \Upsilon_*^i = a_i + b_i(B - V)$. For reference, the mass-to-light ratios predicted by each model for $B - V = 0.6$ are given. The last column denotes whether or not the model includes tracks for TP-AGB stars.

measurements tabulated in the Extragalactic Distance Database (Tully et al. 2009) when available. When no direct distance measurement is known, a Hubble flow distance assuming $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is adopted. Measured colors are given where known. The majority of the sample have observed $B - V$ colors. For brighter galaxies, $J - K$ can readily be extracted from 2MASS (Jarrett et al. 2000). The lower surface brightness galaxies are typically not detected by 2MASS. For many, $V - I$ has been observed (Schombert et al. 2011). References to the sources of these data as well as gas masses and kinematic data are given in the final column of Table 1.

3. STELLAR MASSES FROM POPULATION SYNTHESIS MODELS

Our knowledge of stellar evolution is sufficiently advanced to enable the ab initio calculation of the spectral energy distribution (SED) of complex stellar populations. Considerable effort has gone into the development of stellar population synthesis models that do just this (e.g., Bruzual & Charlot 2003; Maraston 2005). Outstanding progress has been made, and it has become standard practice to quote stellar masses for galaxies based on fits to multi-color data with such models.

The accuracy with which stellar masses can be estimated is debatable, but is probably no better than a factor of two. Outstanding problems include uncertainty in the IMF, variations in the star formation history of galaxies, and the contribution of stars in bright but short-lived phases of evolution (e.g., TP-AGB stars; Maraston 2005). Here we compare the predictions of various models with each other. We also check their internal self-consistency to gage the extent to which the same model predicts the same stellar mass for the same galaxy when the luminosity is measured in different bands.

A simple approach to estimating the stellar mass of a galaxy is to assume a single, constant mass-to-light ratio Υ_* such that $M_* = \Upsilon_* L$. This is a crude approximation, as we expect the mass-to-light ratio of a population to vary with age and, to a lesser extent, with metallicity. For example, using a multi-metallicity model (Schombert & Rakos 2009), we find that a 12 Gyr old stellar population of solar metallicity has $\Upsilon_*^V = 2.8 \text{ M}_\odot/\text{L}_\odot$, while a stellar population with the same age but peak $[\text{Fe}/\text{H}] = -1.5$ has $\Upsilon_*^V = 1.8 \text{ M}_\odot/\text{L}_\odot$. That same solar metallicity stellar population has Υ_*^V of only $0.4 \text{ M}_\odot/\text{L}_\odot$ at an age of 1 Gyr.

Hence a more sophisticated approach is to use a population synthesis model to relate the mean mass-to-light

ratio in band i to a color through

$$\log \Upsilon_*^i = a_i + b_i \text{color}. \quad (1)$$

This reduces to the simple approach if the slope b is small for some band i , as might be expected in the NIR. For the models mentioned above, we find that the solar metallicity model changes in color as it ages from 1 to 12 Gyr by $\Delta(B - V) = 0.37$ and $\Delta(J - K) = 0.03$. We thus expect more variation of Υ_* with optical colors than with those in the NIR, at least for mature stellar populations.

There should be some scatter about the mean Υ_* -color relation. The scatter in Υ_* is also expected to be minimized in the NIR (Bell & de Jong 2001). The scatter in the Υ_* -color relation ultimately limits the accuracy achievable by this approach, even if all else is perfect. One might hope to do better by using multi-color information (e.g., Zibetti et al. 2009), or fitting the entire SED. It is not obvious to what extent additional color information can reduce the scatter in Υ_* given the degeneracy between age and metallicity. In any event, the accuracy of stellar masses inferred from SED fitting is limited by the quality of the population synthesis model that is employed. This approach will obviously be subject to systematic errors if a model differs systematically from reality with wavelength.

The coefficients a_i and b_i are given in Table 2 for several representative models for the $i = V$ and K bands and $B - V$ color. These particular choices are made because the most data are available in these bands. The models typically stop at K while we now have a good deal of Spitzer [3.6] photometry. We relate the population synthesis predicted K -band mass-to-light ratio Υ_*^K to $\Upsilon_*^{[3.6]}$ using the model results of Oh et al. (2008):

$$\Upsilon_*^{[3.6]} = 0.92 \Upsilon_*^K - 0.05. \quad (2)$$

When it is necessary to make this conversion for the data, we assume $K_s - [3.6] = 0.31$. This is the mean color that we obtain for the 74 galaxies of Dale et al. (2005). We assume this value also applies to the many LSB galaxies in our data for which no high quality K -band data are available. Any mean shift in the $K_s - [3.6]$ color between HSB and LSB populations is probably modest, and may well be masked by the scatter, which is only 0.11 mag. in this color in the sample of Dale et al. (2005).

Fig. 1 shows the V and [3.6] band mass-to-light ratios for the models given in Table 2. Rather than simply show a line for each model, we plot the actual galaxy data to emphasize the beads-on-a-string nature of this approach to estimating stellar mass: a single color may provide




Figure 1. The relation between $B - V$ color and the stellar mass-to-light ratio in the V -band (left) and the Spitzer [3.6] band (right) from the population synthesis models of Bell & de Jong (2001) with their scaled Salpeter IMF (open circles), Bell et al. (2003) (filled circles; also scaled Salpeter), Portinari et al. (2004) with a Kroupa (2002) IMF (squares), Zibetti et al. (2009) with a Chabrier (2003) IMF (triangles), and Into & Portinari (2013) with a Kroupa (2002) IMF (stars). The formula of Oh et al. (2008, equation 2) is used to convert between K_s and [3.6]. Note the large disparity between models in the NIR.

a reasonable estimate of the mean mass-to-light ratio, but it cannot reproduce the scatter that one expects. This is estimated to be from 0.1 dex (Bell & de Jong 2001) to 0.15 dex (Portinari et al. 2004) in the K -band, and larger in the optical bands. Even this is an oversimplification, as the scatter may be a function of color, with larger scatter likely in very blue, actively star forming systems. Additionally, the mean color-mass-to-light ratio relation can bend (the slope becomes much steeper for $B - V < 0.55$ in the models of Portinari et al. 2004) or even bifurcate: the line of Bell et al. (2003) splits the difference between distinct branches of high and low Υ_*^K at blue colors in their Fig. 20.

Different models will differ if they adopt different evolutionary tracks or a different IMF. Both matter for the models considered here. As pointed out by Bell & de Jong (2001), changes in the IMF to include more or fewer low mass stars serve mostly to change the mass without much altering the luminosity or color, so to a decent approximation can be treated as shifts in Υ_* . For specificity, we adopt the scaled Salpeter IMF of Bell & de Jong (2001) for their model and also for that of Bell et al. (2003). We adopt the Kroupa IMF (Kroupa 2002) for the model of Portinari et al. (2004) and Into & Portinari (2013), while Zibetti et al. (2009) uses the Chabrier IMF (Chabrier 2003). These models all give a similar run of Υ_* with color in the optical (left panel of Fig. 1), with small offsets for the different IMFs as well as other detailed differences. The agreement is less good in the NIR (right panel of Fig. 1). There is a huge disparity between the model of Bell et al. (2003), which has a relatively flat slope b_k with a high normalization $\Upsilon_*^{[3.6]} > 0.55 M_\odot/L_\odot$, and that of Zibetti et al. (2009), which has a steep dependence on color (even in the NIR) and a low normalization $\Upsilon_*^{[3.6]} < 0.3 M_\odot/L_\odot$. The model of Into & Portinari (2013) is intermediate between that of Zibetti et al. (2009) and the older models. The models of Bell & de Jong (2001) and Portinari et al. (2004) largely overlap in between these two extremes, but it is unclear whether we should take comfort in this apparent agreement as the two have different IMFs. Bell & de Jong (2001) give a 0.15 dex offset between

their scaled Salpeter IMF and the Kroupa IMF, in the sense that the latter is lighter. However, the Kroupa case appears not to include brown dwarfs (Bell et al. 2003), which would increase the mass by about 0.15 dex. Here we simply adopt the various published coefficients without trying to reconcile definitions of the IMF.

The chief difference in evolutionary tracks between the various models considered here is the inclusion of TP-AGB stars by Zibetti et al. (2009) and Into & Portinari (2013) using the models of Maraston (2005). These stars are in the latest stages of evolution, being short-lived and rare, but quite bright. Their contribution to the integrated luminosity of stellar populations is most pronounced in the NIR, where they greatly enhance the predicted luminosity of galaxies while doing little to alter the predictions of previous generations of models in the optical portion of the spectrum. This results in the low mass-to-light ratios of the models¹ of Zibetti et al. (2009) and Into & Portinari (2013) in the right hand panel of Fig. 1.

We use the models in Table 2 with the data in Table 1 to compute the stellar masses of sample galaxies. These are reported in Table 3. For each galaxy, we use the observed color to predict the mass-to-light ratio separately in both the optical and NIR for each model. We then use the corresponding luminosity to obtain a stellar mass estimate. The models of Bell & de Jong (2001) and Portinari et al. (2004) are barely distinguishable, so we tabulate only the latter. This results in eight distinct mass estimates for each galaxy: two for each of the four models. For a few galaxies where only $V - I$ is available and not $B - V$, we utilize the available color with the same parameters as the other models, when possible (Bell et al. (2003) and Zibetti et al. (2009) do not tabulate coefficients for $V - I$).

We can now compare stellar mass estimate from different population synthesis models. We can also check each model for internal self-consistency. If all is well, the stellar mass estimated for the same galaxy will be the same

¹ Zibetti et al. (2009) advocate using multiband colors to estimate mass-to-light ratios, and their single-color coefficients that are reproduced in Table 2 are only approximations made to facilitate the sort of comparison made here.

Table 3
Stellar Masses from Population Synthesis Models

Galaxy	B03		P04		Z09		IP13	
	M_*^V	$M_*^{[3.6]}$	M_*^V	$M_*^{[3.6]}$	M_*^V	$M_*^{[3.6]}$	M_*^V	$M_*^{[3.6]}$
D564-8	...	...	7.14	6.99	...	...	7.08	6.86
DDO 154	7.54	7.63	7.51	7.24	7.27	6.45	7.37	7.01
D631-7	7.64	7.77	7.61	7.45	7.41	6.80	7.50	7.26
D640-13	...	...	7.20	7.21	...	...	6.95	6.69
D568-2	7.73	7.81	7.70	7.51	7.52	6.90	7.60	7.35
D572-5	7.70	7.81	7.67	7.51	7.49	6.89	7.57	7.34
F415-3	8.19	7.90	8.16	7.72	8.08	7.27	8.12	7.63
D500-3	...	...	7.35	7.03	...	...	...	...
DDO 168	8.00	8.04	7.97	7.66	7.72	6.87	7.83	7.43
D575-1	...	...	7.39	7.68	...	...	7.21	7.37
D512-2	...	...	7.96	7.92	...	...	7.83	7.70
F611-1	8.21	8.27	8.17	8.06	8.06	7.56	8.12	7.94
D500-2	8.53	8.39	8.50	8.15	8.36	7.61	8.43	8.01
F565-V2	8.36	8.58	8.33	8.28	8.15	7.67	8.23	8.11
NGC 2366	9.09	8.65	9.06	8.42	8.93	7.91	9.00	8.30
D723-5	8.78	8.78	8.75	8.56	8.63	8.05	8.69	8.43
F563-V1	8.48	8.96	8.45	8.51	8.16	7.51	8.29	8.23
IC 2574	9.26	9.14	9.23	8.82	9.04	8.18	9.13	8.64
F563-1	8.95	9.24	8.91	8.91	8.71	8.24	8.80	8.72
F574-2	9.38	9.30	9.35	9.10	9.24	8.61	9.30	8.99
NGC 2976	9.35	9.30	9.32	9.08	9.20	8.57	9.26	8.96
F568-V1	9.36	9.41	9.32	9.14	9.16	8.55	9.22	8.98
F561-1	9.52	9.47	9.49	9.34	9.44	8.94	9.47	9.27
F577-V1	9.44	9.47	9.40	9.21	9.26	8.66	9.33	9.07
NGC 1003	10.00	9.52	9.97	9.21	9.78	8.57	9.87	9.03
UGC 5005	9.24	9.54	9.21	9.18	8.98	8.44	9.08	8.96
F574-1	9.61	9.67	9.58	9.39	9.43	8.84	9.50	9.25
F568-1	9.54	9.65	9.51	9.40	9.37	8.86	9.44	9.27
NGC 7793	9.89	9.69	9.86	9.52	9.78	9.08	9.83	9.43
UGC 128	9.85	9.86	9.81	9.69	9.73	9.24	9.78	9.60
NGC 2403	9.74	9.86	9.71	9.53	9.50	8.85	9.59	9.33
NGC 925	10.17	10.01	10.14	9.75	9.99	9.20	10.06	9.61
NGC 2903	10.61	10.19	10.58	9.97	10.46	9.46	10.52	9.85
NGC 3198	10.34	10.28	10.30	9.97	10.12	9.34	10.20	9.80
NGC 3621	10.21	10.30	10.18	10.06	10.04	9.53	10.11	9.93
NGC 3521	10.84	10.79	10.80	10.66	10.75	10.24	10.79	10.58
NGC 3031	10.99	10.85	10.96	10.80	10.98	10.48	10.99	10.78
NGC 5055	10.84	10.95	10.81	10.79	10.74	10.35	10.78	10.70
NGC 2998	10.83	10.99	10.80	10.70	10.63	10.09	10.71	10.53
NGC 2841	11.05	11.07	11.01	10.98	11.00	10.61	11.02	10.93
NGC 6674	10.92	11.17	10.88	10.96	10.77	10.47	10.83	10.85
NGC 7331	11.05	11.23	11.01	11.06	10.93	10.61	10.98	10.97
NGC 801	11.02	11.24	10.99	11.05	10.90	10.59	10.95	10.96
NGC 5533	11.17	11.32	11.14	11.24	11.14	10.89	11.15	11.20
UGC 2885	11.24	11.46	11.20	11.18	11.04	10.60	11.11	11.03

Note. — Masses are base ten logarithms in $M_\odot$. For each population model, the mass estimate from the V -band luminosity is given first, then that from the $[3.6]$ luminosity. The models used are those of Bell et al. (2003), Portinari et al. (2004), Zibetti et al. (2009), and Into & Portinari (2013).

irrespective of whether the luminosity is measured in the optical or NIR. Scatter in the color–mass-to-light relation precludes this from ever being exactly true, but the sample is large enough that we can check whether it is true on average. This is done in Fig. 2, which shows the stellar mass estimated from the $[3.6]$ luminosity plotted against that estimated from the V -band luminosity.

The external consistency of the models is fairly good in the optical: examination of Table 3 shows that M_*^V is usually similar across the board, with small offsets that are roughly consistent with differences in the adopted IMFs and the intrinsic uncertainties of complex stellar populations. The same cannot be said in the NIR. Indeed, most of the models are not internally self-consistent: for a given stellar mass, they over-predict the infrared luminosity relative to the optical luminosity. In other words, the predicted infrared mass-to-light ratios

are too small relative to those in the optical.

The data cover many decades in luminosity. Consequently, even a small offset from the line of equality in the logarithmic Fig. 2 corresponds to a serious misestimation of the mass-to-light ratio. The models of Bell & de Jong (2001) and Portinari et al. (2004) suffer this problem, but it becomes more severe in the more recent models of Zibetti et al. (2009) and Into & Portinari (2013). In the case of Zibetti et al. (2009), the offset in mass between optical and NIR estimates is typically greater than half a dex.

The chief difference between the older and newer models is the inclusion of TP-AGB stars in the more recent models. While it is obviously desirable to include all the relevant evolutionary phases, it appears that the contribution of the TP-AGB population to the integrated luminosity of galaxies is overstated by the models of




Figure 2. Stellar masses (Table 3) estimated from luminosities and colors using the formulae supplied by population synthesis models (Bell et al. 2003; Portinari et al. 2004; Zibetti et al. 2009; Into & Portinari 2013). For each model, the mass estimated from the [3.6] luminosity of each galaxy is plotted against that estimated from the V-band luminosity. Filled circles use $B - V$ colors to estimate mass-to-light ratios; open circles use $V - I$ colors. If the models were perfect the data would follow the line of unity, modulo the expected scatter in the relation between the mass-to-light ratio and color. With the exception of the model of Bell et al. (2003), the models tend to over-predict the NIR luminosity relative to the optical luminosity for a given stellar mass.

Maraston (2005). This conclusion is consistent with the finding of Kriek et al. (2010), who attempted to fit these models to the SEDs of post-starburst galaxies. Similarly, Zibetti et al. (2013) do not observe the spectral features expected from TP-AGB stars in the NIR spectra of post-starburst galaxies. Melbourne et al. (2012) also reach a similar conclusion about the ratio of NIR to optical luminosity from the color-magnitude diagrams of nearby galaxies.

Our understanding of the latest phases of stellar evolution does not yet appear to be sufficiently complete to confidently predict the NIR spectra of complex stellar populations. All *ab initio* models considered here suffer the same problem, which has gotten worse rather than better with good faith efforts to extend evolutionary tracks to late times. Our results provide an empirical constraint to help inform future modeling. In the mean time, considerable caution is warranted in assigning stellar masses based on SED fits (Conroy & Gunn 2010). The results will depend not only on the model adopted, but also on what range of wavelength is fit. The offset

between optical and NIR luminosity identified here will result in a systematic skew towards lower stellar masses as more NIR data are incorporated into SED fits.

The model of Bell et al. (2003) is not entirely *ab initio*, as it makes extensive use of empirical constraints. This grounding in data makes it the only model to come close to the ideal of obtaining equal masses independently from optical and NIR luminosities. This might be a bit of a fluke given the bifurcation noted above, but nevertheless the ratio of optical to NIR flux is approximately that observed in real galaxies.

Referring back to Fig. 1, we can immediately take two lessons: (i) self-consistency is only obtained for relatively high NIR mass-to-light ratios, and (ii) the slope of the color-mass-to-light ratio is not steep in the NIR. Consistency over the full range of the data only follows if $\Upsilon_*^{[3.6]}$ is approximately constant. It has, of course, been suspected for a long time that Υ_* would be roughly constant in the NIR. The data presented here support that expectation.

We can perform a simple exercise to pursue the appar-




Figure 3. Stellar masses as in Fig. 2 but assuming a constant mass-to-light ratio at $[3.6]$ rather than the population synthesis value. The values used are those that on average agree with the V -band stellar mass: $\Upsilon_*^{[3.6]} = 0.58; 0.53; 0.43;$ and $0.48 \text{ M}_\odot/\text{L}_\odot$ for the models of Bell et al. (2003); Portinari et al. (2004); Zibetti et al. (2009); and Into & Portinari (2013), respectively. Comparison to Fig. 2 makes it clear that a constant mass-to-light ratio in the NIR provides a more self-consistent description of the data than do models that apply a color correction in that part of the spectrum.

ent constancy of $\Upsilon_*^{[3.6]}$. In Fig. 3 we retain the stellar mass predicted from the optical luminosity (M_*^V from Table 3), but for the NIR based mass we adopt a constant mass-to-light ratio for all galaxies. We then ask what value of $\Upsilon_*^{[3.6]}$ best reconciles the data with the optically estimated mass. The best fit mean NIR mass-to-light ratio is $\Upsilon_*^{[3.6]} = 0.52; 0.58; 0.53; 0.43;$ and $0.48 \text{ M}_\odot/\text{L}_\odot$ for the models of Bell & de Jong (2001); Bell et al. (2003); Portinari et al. (2004); Zibetti et al. (2009); and Into & Portinari (2013), respectively. Note that $\Upsilon_*^{[3.6]}$ must be relatively large to obtain self-consistency even for the model with the lowest mean mass-to-light ratio (Zibetti et al. 2009).

Fig. 3 now looks like it should: a one to one correspondence between stellar mass estimated from optical and NIR luminosity, with the inevitable scatter in Υ_* from galaxy to galaxy. Simply assuming a constant $\Upsilon_*^{[3.6]}$ performs better than the models considered here (compare Figures 2 and 3). Only in the case of Bell et al. (2003) are the two not statistically distinguishable. As noted above, the relative success of the Bell et al. (2003) model

stems in part from its relatively flat slope, $b_K = 0.135$. All of the other models listed in Table 3 have $b_K > 0.65$, and two have $b_K > 1$. Such steep slopes prevent self-consistency. There is no strong evidence in the data for a non-zero slope in the color-mass-to-light relation at $3.6\mu\text{m}$. A small slope is of course allowed, but may well be lost in the scatter in Υ_* .

The absolute value of the required $\Upsilon_*^{[3.6]}$ does not vary much from model to model. Given that these models bracket the plausible range in IMFs, there is tentative reason to hope that the absolute value of $\Upsilon_*^{[3.6]}$ estimated in this way is approximately correct. Apparently, $\Upsilon_*^{[3.6]} \approx 0.5 \text{ M}_\odot/\text{L}_\odot$ to a good approximation, with a range of about 20% from the Chabrier IMF at the low end to scaled Salpeter at the high end.

The absolute value of the mass-to-light ratio given here implicitly assumes that the modeling of the optical portion of the spectrum is essentially correct. Really the data only constrain the ratio of $\Upsilon_*^{[3.6]} : \Upsilon_*^V$ to be higher than most models indicate. One is free to adjust the normalization of both optical and NIR mass-to-light ratio so




Figure 4. Left panel: the stellar mass estimated from the Baryonic Tully-Fisher Relation (Table 4) as a function of V -band luminosity. Right panel: the BTFR stellar mass as a function of $[3.6]$ luminosity. Downward pointing triangles represent 2σ upper limits for galaxies with stellar masses that are formally negative. The lines are the lines of equality for $\Upsilon_*^V = 1.3$ and $\Upsilon_*^{[3.6]} = 0.5 \text{ M}_\odot/\text{L}_\odot$. Luminosity is a good proxy for kinematic stellar mass, with somewhat less scatter in the NIR than in the optical.

long as the appropriate ratio is preserved. However, obtaining a NIR mass-to-light ratio much outside the range $0.4 < \Upsilon_*^{[3.6]} < 0.6 \text{ M}_\odot/\text{L}_\odot$ implies an IMF² outside the bounds of recent determinations (Kroupa 2002).

4. STELLAR MASSES FROM THE BARYONIC TULLY-FISHER RELATION

An independent method to estimate stellar masses is provided by the Baryonic Tully-Fisher Relation (BTFR). The observed mass of baryons — stars plus gas — is observed to correlate well with rotation velocity (McGaugh et al. 2000; Verheijen 2001; Bell & de Jong 2001; McGaugh 2005). Recent work on gas dominated galaxies (Begum et al. 2008; Stark et al. 2009; Trachternach et al. 2009) has made it possible to obtain an absolute calibration of the mass scale of the BTFR that is largely independent of stellar mass estimates (Stark et al. 2009; McGaugh 2011). This provides a powerful new constraint on stellar masses.

The BTFR can be written as

$$M_b = AV_f^4 \quad (3)$$

where V_f is the rotation velocity measured in the flat outer portion of a resolved rotation curve and the baryonic mass $M_b = M_* + M_g$. A recent calibration of the BTFR obtains $A = 47 \pm 6 \text{ M}_\odot \text{ km}^{-4} \text{ s}^4$ (McGaugh 2012). The gas masses in this calibration were taken to be $M_g = 1.33M_{HI}$, so the same is done here for consistency.³

² Our experience with point source contamination in the Spitzer data confirms that there are indeed stars that only become noticeable in the NIR. However, these apparently are foreground M dwarfs, the profusion of which anecdotally suggests that the IMF is not lacking in low mass stars.

³ A few of the sample galaxies have molecular gas masses estimated from CO observations (Leroy et al. 2008). We have experimented with including molecular gas from this and from estimates based on the observed star formation rate. Unfortunately, such data do not exist for all sample galaxies. For those that do, the result is essentially unchanged if we allow for an appropriately higher calibration of the BTFR ($A = 50 \pm 6 \text{ M}_\odot \text{ km}^{-4} \text{ s}^4$; McGaugh 2005). The stellar masses of individual galaxies change slightly, but the mean and scatter in the mass-to-light ratios are indistinguishable.

In order to estimate stellar masses from the BTFR, we use the measured rotation velocity and equation 3 to first estimate the baryonic mass.⁴ The stellar mass then follows by subtracting the gas mass:

$$M_* = M_b - M_g = AV_f^4 - 1.33M_{HI}. \quad (4)$$

The data necessary to make this computation are the rotation velocity and the mass of atomic hydrogen. These are taken from various sources: de Blok et al. (1996); McGaugh (2005); Begum et al. (2008); de Blok et al. (2008); Swaters et al. (2009); Stark et al. (2009); Trachternach et al. (2009) and compiled in Table 4. The most restrictive requirement on the data is the need for a resolved rotation curve; this does not exist for all galaxies in Table 1, but we have succeeded in collecting a sample that covers a large dynamic range in V_f as well as in luminosity.

The results of the computation of equation 4 are given in Table 4 as stellar mass and the mass-to-light ratios Υ_*^V and $\Upsilon_*^{[3.6]}$. These are shown in Fig. 4. The stellar mass estimated in this way from the BTFR correlates as expected with luminosity, especially $L_{[3.6]}$, even though equation 4 involves only rotation velocity and gas mass, and makes no reference to the observed luminosity. Furthermore, the mean normalization is consistent with $\Upsilon_*^{[3.6]} = 0.5 \text{ M}_\odot/\text{L}_\odot$. This provides a kinematic constraint on Υ_* that is entirely consistent with what we inferred in the previous section. Stellar mass estimates from the BTFR are consistent with but independent of those based on requiring self-consistency in population synthesis.

It is satisfying that the two independent methods for estimating stellar mass, population synthesis and the BTFR, provide convergent results. It is worth noting at this juncture a few exceptions that prove the rule. Looking at equation 4, it is remarkable that this works as well as it does. Starting from V_f which is typically of order

⁴ The empirical BTFR involves only the observed baryonic mass and the rotation velocity, and does not directly involve the mass of dark matter even if it is indirectly implicated in setting the amplitude of the velocity. See McGaugh (2011, 2012) for further discussion.

Table 4
Stellar Masses from the BTFR

Galaxy	V_f	M_g	M_*	σ_{M_*}	Υ_*^V	$\sigma_{\Upsilon_*^V}$	$\Upsilon_*^{[3.6]}$	$\sigma_{\Upsilon_*^{[3.6]}}$
D564-8	29	0.0210	0.0126	0.0228	1.30	2.34	0.56	1.00
KK98 251	40	0.123	-0.0027	0.0392	...	...	-0.04	0.55
DDO 154	53	0.420	-0.0410	0.142	-0.73	2.51	-0.56	1.95
D631-7	53	0.195	0.172	0.145	3.20	2.69	1.76	1.48
D500-3	45	0.0874	0.112	0.107	0.66	0.63	0.76	0.73
DDO 168	53	0.421	-0.0394	0.0866	-0.24	0.53	-0.21	0.46
D512-2	38	0.0926	0.0003	0.0659	0.0032	0.63	0.0012	0.24
D500-2	68	1.15	-0.147	0.415	-0.48	1.36	-0.37	1.04
DDO 170	65	1.14	-0.302	0.149	...	...	-0.61	0.30
NGC 2366	60	0.798	-0.201	0.419	-0.19	0.40	-0.28	0.58
IC 2574	77	1.88	-0.249	0.434	-0.11	0.20	-0.11	0.19
F563-1	111	4.65	2.48	2.72	2.20	2.41	0.86	0.95
NGC 2976	86	0.178	2.39	0.531	1.30	0.29	0.75	0.17
F568-V1	124	5.22	5.89	3.85	2.50	1.63	1.39	0.91
NGC 1003	114	6.78	1.01	1.12	0.084	0.093	0.18	0.20
UGC 5005	99	7.04	-2.52	1.91	-0.98	0.74	-0.43	0.33
F574-1	100	5.46	-0.766	1.97	-0.21	0.53	-0.11	0.28
F568-1	116	5.34	3.16	3.12	1.02	1.00	0.45	0.44
NGC 7793	110	1.01	5.97	1.42	1.18	0.28	0.78	0.19
UGC 128	131	8.66	5.177	4.58	1.15	1.02	0.46	0.41
NGC 2403	136	3.35	12.6	2.77	1.74	0.38	1.03	0.23
NGC 925	114	5.31	2.52	2.00	0.18	0.14	0.15	0.12
NGC 2903	188	5.78	52.3	10.3	1.56	0.31	2.10	0.42
NGC 3198	149	13.5	9.57	3.93	0.38	0.16	0.31	0.13
NGC 3621	152	9.28	16.0	3.86	1.11	0.27	0.49	0.12
NGC 3521	192	5.96	57.8	24.7	1.52	0.65	0.61	0.26
NGC 3031	199	9.27	64.4	22.5	1.80	0.63	0.63	0.22
NGC 5055	181	9.59	41.0	14.7	0.95	0.34	0.30	0.11
NGC 2998	212	36.8	57.5	12.9	0.77	0.17	0.36	0.08
NGC 2841	279	11.4	272.	58.8	5.28	1.14	1.52	0.33
UGC 2953	260	9.28	205.	59.6	...	...	0.97	0.28
NGC 6674	241	50.8	106.	22.8	1.68	0.36	0.46	0.10
NGC 7331	246	12.5	159.	32.0	2.24	0.45	0.61	0.12
NGC 801	219	48.8	58.9	15.3	0.83	0.22	0.22	0.06
NGC 5533	240	31.2	124.	23.7	1.98	0.38	0.41	0.08
UGC11455	272	24.9	232	42.2	...	...	0.67	0.12
UGCA145	300	52.1	329.	54.8	...	...	0.94	0.16
UGC 2885	298	45.8	324.	68.6	1.82	0.39	0.69	0.15

Note. — Stellar masses and mass-to-light ratios inferred from the Baryonic Tully-Fisher relation (McGaugh 2012). The rotation velocity is in km s^{-1} . Masses are in units of $10^9 M_\odot$. The gas mass assumes $M_g = 1.33 M_{HI}$. Mass-to-light ratios are in solar units $M_\odot/L_\odot$.

$\sim 100 \text{ km s}^{-1}$, we first raise this number to the fourth power. We then multiply by an empirically determined constant of the same order. This gives a large number appropriate to the baryonic mass of a galaxy. We then subtract another large number, the gas mass. Anyone with experience in attempting this sort of calculation with astronomical data will immediately recognize the dangers inherent in such an approach. Raising a number to a large power invites the propagation of errors; subtracting one large number from another practically begs them to blow up. Remarkably, this rarely happens. This is a credit to both the quality of the data and the virtue of the method.

It does sometimes happen that the stellar mass estimate from equation 4 is negative. This is especially prone to occur in gas rich galaxies where the gas mass accounts for nearly all of the baryonic mass. There are nine occurrences of this among the 38 galaxies listed in Table 4. The formal significance of these negative masses is $\lesssim 2\sigma$ in all cases, so there are no examples where the method yields a definitively unphysical result. Indeed, it would be worrisome if one did not obtain the occasional nega-

tive mass⁵, especially given the large number of gas rich galaxies in this sample (about half have $f_g > 0.5$).

At the opposite extreme, maximum disk places an upper limit on the allowed stellar mass: the inferred mass should not cause the rotation curve of the stellar component to exceed the observed rotation curve. Since the BTFR only unitizes the outer part of the rotation curve, it is in principle quite possible for this to occur. In this sample, it occurs exactly once: in the case of NGC 2841. This is hardly surprising, as NGC 2841 is a well known outlier from the Tully-Fisher relation (Bottema et al. 2002). If it is a true outlier, then this method does not apply. It remains conceivable that its distance is wrong or some other systematic error is at fault. For example, an undiagnosed warp in the outer regions could cause an error in V_f . Regardless of what is really going on in this case, the mass returned by the BTFR is unphysical in that it exceeds maximum disk.

⁵ We have attempted this exercise using line-widths instead of V_f from rotation curves. Line-widths fail by giving many negative stellar masses and some unreasonably large Υ_* . This happens because of the larger intrinsic scatter of the line-width based BTFR (McGaugh 2012): good, resolved rotation curve are essential for estimating Υ_* .




Figure 5. Mass-to-light ratios inferred from the baryonic Tully-Fisher relation in the V -band (top) and at $[3.6]$ (bottom) as a function of $B - V$ (left) and $J - K_s$ (right) color. Galaxies for which the uncertainty in the stellar mass exceeds half of the stellar mass are shown as smaller symbols. Downward pointing triangles represent 2σ upper limits for galaxies with formally negative Υ_* . NGC 2841 is shown as an open circle as the inferred stellar mass exceeds its maximum disk value. For comparison, the stellar population models of Bell et al. (2003, solid lines) and Portinari et al. (2004, dashed lines) are shown in the left panels. The median mass-to-light ratios are $\Upsilon_*^V = 1.09 M_\odot/L_\odot$ and $\Upsilon_*^{[3.6]} = 0.45 M_\odot/L_\odot$ (dotted lines in right panels). For the average $K_s - [3.6] = 0.31$ color, the corresponding K_s -band mass-to-light ratio is $\Upsilon_*^{K_s} = 0.58 M_\odot/L_\odot$. Note that the majority of galaxies are within 2σ of the median value in the NIR.

The only other galaxy that appears troublesome is NGC 2903. Its stellar mass is not unphysical, but it is odd in having its BTFR-inferred NIR mass-to-light ratio larger than the optical value: $\Upsilon_*^{[3.6]} > \Upsilon_*^V$. Population synthesis models do not anticipate that this should happen, even after the upward correction to $\Upsilon_*^{[3.6]}$ discussed in §3, nor does it: all other galaxies but two⁶ have BTFR-inferred $\Upsilon_*^{[3.6]} < \Upsilon_*^V$. Looking at Table 3, the stellar masses for NGC 2903 from optical and NIR are badly offset according to all four models, as if the NIR luminosity had been underestimated. Indeed, from a purely empirical perspective, NGC 2903 is an outlier from the infrared luminosity Tully-Fisher relation even

though it is not from the optical version of the relation. The source of the photometry is de Blok et al. (2008), who note problems in constructing the mass model for this galaxy, so we regard its $[3.6]$ luminosity as dubious.

This is a small list of exceptional cases for a method that should be so easily derailed. Just as the stellar mass might come out negative, it could also come out an order of magnitude or more too heavy. That this does not often happen is encouraging.

Fig. 5 shows the mass-to-light ratios inferred from the BTFR as a function of color. The data show a trend of Υ_*^V increasing with $B - V$ color that is consistent with the expectations of population models. This has already been seen to be the case in the B -band (Stark et al. 2009); the present data are entirely consistent with that earlier result. There is a fair amount of scatter in Υ_* , but this is also expected. The inferred optical mass-to-light ratios behave as they should from the perspective of population synthesis.

There is no clear indication of a correlation of either

⁶ D500-3 and NGC 1003 have $\Upsilon_*^{[3.6]} \approx \Upsilon_*^V$, indistinguishable within the errors. More generally, population models do anticipate low and rapidly variable mass-to-light ratios for populations dominated by very young stars, but this situation stabilizes to $\Upsilon_*^{[3.6]} < \Upsilon_*^V$ for most relevant ages (> 1 Gyr).




Figure 6. Mass-to-light ratios inferred from the baryonic Tully-Fisher relation as per Fig. 5, but restricted to the more precise data with $\sigma_{M_*} < \frac{1}{2} M_*$. The mean mass-to-light ratios are $\Upsilon_*^V = 1.3 M_\odot/L_\odot$ and $\Upsilon_*^{[3.6]} = 0.5 M_\odot/L_\odot$ (dotted line in right panel). For the average $K_s - [3.6]$ color, the corresponding K_s -band mass-to-light ratio is $\Upsilon_*^{K_s} = 0.65 M_\odot/L_\odot$. This is slightly heavier than seen for all the data in Fig. 5 as the more precise data tend to be for the brighter, more evolved spirals. The line in the left panel represents a fit to the data with the slope fixed to the value anticipated by population synthesis models (Table 5).

mass-to-light ratio with $J - K_s$ color. This is not surprising, since this color should be more sensitive to the metallicity of stars than their age (Bell & de Jong 2000). This means that NIR colors are a poor indicator of Υ_* . One is better off assuming constant $\Upsilon_*^{[3.6]}$ than attempting to correct Υ_* with a color term involving NIR colors.

The NIR mass-to-light ratio is also insensitive to optical color. The data allow for a small slope $b_{[3.6]}$, but are also consistent with zero slope. This is consistent with the results of §3, where it was seen that small slopes were required for population synthesis models to be internally self-consistent. To a reasonable approximation, $\Upsilon_*^{[3.6]}$ is constant. The median mass-to-light ratio is $\Upsilon_*^{[3.6]} = 0.45 M_\odot/L_\odot$. For the average $K_s - [3.6] = 0.31$ color, the corresponding mass-to-light ratio is $\Upsilon_*^{K_s} = 0.58 M_\odot/L_\odot$.

The scatter in Fig. 5 is somewhat misleading because of the large uncertainty in stellar mass estimated from the BTFR for gas rich galaxies. In Fig. 6 we restrict the sample to those galaxies with formal errors in mass $\sigma_{M_*}/M_* < 50\%$. We also exclude NGC 2841, whose BTFR mass is unphysical, and NGC 2903, whose [3.6] luminosity we consider untrustworthy (see above). This leaves a “clean” sub-sample of 14 galaxies.

The scatter in Fig. 6 is reduced from that in Fig. 5, as expected by construction. The V-band mass to-light-ratios remain consistent with the color trend expected from population models, but there remains some scatter. The [3.6] mass-to-light ratio is effectively constant. There is no hint of a trend with color, and many individual galaxies are consistent with the average value within the errors. The average mass-to-light ratio is slightly higher for the clean sub-sample, with a mean $\Upsilon_*^{[3.6]} = 0.5 M_\odot/L_\odot$ (equivalent to $\Upsilon_*^{K_s} = 0.65 M_\odot/L_\odot$). This small shift from $0.45 M_\odot/L_\odot$ for the entire sample is too small to consider significant. Nevertheless, it goes in an intriguing direction, hinting that gas rich galaxies may tend to have lower mass-to-light ratios, even in the NIR. It is not difficult to imagine that they have younger ages and/or are more metal poor than bright, well-developed spirals.

The clean sub-sample gives us some handle on the intrinsic scatter in mass-to-light ratios. The scatter in $\Upsilon_*^{[3.6]}$ is 0.23 dex while the average uncertainty is 0.19 dex. This implies an intrinsic scatter of 0.13 dex. While the sample is too small and the uncertainty in the uncertainties too great to draw firm conclusions from this, this implied value of the intrinsic scatter is nicely bracketed by that anticipated in the NIR by the models of Bell & de Jong (2001, 0.1 dex) and Portinari et al. (2004, 0.15 dex).

Table 5 summarizes our calibration of the color—mass-to-light ratio relation determined from the BTFR. This is entirely consistent with the expectation of population synthesis models once those are made self-consistent as discussed in §3. We do not have enough data to constrain the slope of the relation, which we fix to a representative value from the population models in the optical bands. The B -band calibration is given along with that in the V -band for completeness; see also Stark et al. (2009). In the NIR there is no indication of a systematic variation of Υ_* with color, so we fix the slope to zero. The best fit to the NIR Υ_* is a constant, differing slightly by sample selection. The clean sub-sample gives a more precise constraint, but may build in a small systematic bias: the uncertainty on the stellar mass from the BTFR is necessarily smaller for star dominated rather than gas dominated galaxies. However, the difference is not large ($\Upsilon_*^{[3.6]} = 0.5$ versus $0.45 M_\odot/L_\odot$), and we do not consider it to be meaningful.

The basic value of the BTFR constrained mass-to-light ratio is meaningful. As discussed by Stark et al. (2009), it provides a constraint on the IMF. Comparing a_V determined from the BTFR in Table 5 with that expected from population synthesis models in Table 2, it appears that the data are most consistent with a Kroupa IMF (Kroupa 2002).

5. DISCUSSION

For all intents and purposes, the data appear to be consistent with a nearly constant NIR mass-to-light ratio $\Upsilon_*^{[3.6]} = 0.5 M_\odot/L_\odot$. This value is inferred in two

Table 5
Self-Consistent Stellar Mass-to-Light Ratios

Sample	a_B	b_B	a_V	b_V	a_{K_s}	b_{K_s}	$a_{[3.6]}$	$b_{[3.6]}$	$\Upsilon_{0.6}^B$	$\Upsilon_{0.6}^V$	$\Upsilon_{0.6}^{K_s}$	$\Upsilon_{0.6}^{[3.6]}$
All galaxies	-0.94	1.7	-0.68	1.3	-0.24	0	-0.35	0	1.20	1.26	0.58	0.45
Clean sub-sample	-0.90	1.7	-0.64	1.3	-0.19	0	-0.30	0	1.32	1.38	0.65	0.50

Note. — Stellar mass-to-light ratios (in solar units) $\log \Upsilon_*^i = a_i + b_i(B - V)$ tied to the gas rich galaxy calibration of the BTFR (McGaugh 2012). The slope is fixed so that only the zero point varies with sample. The clean sub-sample consists of galaxies with relatively precise stellar masses estimated from the BTFR with formal uncertainties $\sigma_{M_*} < \frac{1}{2} M_*$. Implied mass-to-light ratios at $B - V = 0.6$ are given for filters $i = B, V, K_s$, and $[3.6]$.

independent ways: (i) by requiring self-consistency in population synthesis models, and (ii) from the stellar mass required to obey the BTFR fixed by gas dominated galaxies. The equivalent K_s -band mass-to-light ratio is $\Upsilon_*^K = 0.65 M_\odot/L_\odot$ assuming $K_s - [3.6] = 0.31$ with solar absolute magnitudes $M_\odot^{K_s} = 3.27$ and $M_\odot^{[3.6]} = 3.24$ (Oh et al. 2008).

The apparent near-constancy of the stellar mass-to-light ratio in the NIR that we infer here has long been anticipated. This observation has important ramifications. Simply by observing the NIR luminosity of a galaxy, we obtain a good handle on its stellar mass — no further input is required.

It is tempting to conclude not only that the NIR luminosity of a galaxy is a good indicator of its stellar mass, but also that we have constrained the conversion of light to mass to within $\sim 20\%$ of $\Upsilon_*^{[3.6]} = 0.5 M_\odot/L_\odot$. There is certainly real scatter within that range, and there are hints in the data that the range $0.4 - 0.5$ may be more likely among gas rich disks than the range $0.5 - 0.6$. While it is easy to imagine that some galaxies with genuinely young stellar populations have even lower Υ_* , this does not appear to be the general rule.

Our chief hesitation in declaring that the problem of stellar mass is solved (besides historical precedence that such assertions are never warranted) is that the DiskMass survey (Bershady et al. 2010) finds masses that appear to be systematically lower (Bershady et al. 2011) by a factor of two or even a bit more. There is no overlap in the samples, so this estimate is inferred from their estimate of the maximality of disks, which is small, whereas our Υ_* are more in line with, or even a bit heavier than those of de Blok et al. (2008, see their Figure 61). We do not know the reason for this inconsistency.

We can note some consequences for our study of a much lower mass-to-light ratio. From the stellar population perspective, it would be possible, at least in principle, to reduce Υ_* of both optical and NIR bands while preserving the ratio found in §3. In practice this is difficult. The optical portion of the spectrum is reasonably well understood (e.g., Kriek et al. 2010), so such a reduction would correspond to a large shift in the IMF. That is not inconceivable, but would have real consequences for star counts: we cannot reduce the IMF arbitrarily and still account for the stars we see directly.

It is also difficult to reconcile lower Υ_* with the BTFR. The observed slope of the K -band Tully-Fisher relation (Verheijen 2001) is 4, consistent with that of gas dominated galaxies (McGaugh 2012). The two join continuously if and only if the stellar mass-to-light ratio is as

we estimate. If a lower mass-to-light ratio is adopted, the BTFR develops a break in its slope and the scatter increases (McGaugh 2005). There is no hint of a break in the empirical relation, so it would be odd to insert one. It would also be odd if the scatter in the mass-based BTFR were *larger* than that in the luminosity-based Tully-Fisher relation, since variations in Υ_* should only add scatter when going from the intrinsic BTFR to the luminous Tully-Fisher relation. This would be especially strange since the expected increase in scatter due to Υ_* as one goes from the NIR to progressively bluer bands is observed (Verheijen 2001). We therefore see no way to gracefully reconcile our results with those of Bershady et al. (2011), so there appears to remain of systematic uncertainty in stellar mass of a factor of ~ 2 .

5.1. Gas Fractions of Disk Galaxies

Gas fractions provide an important clue to the evolutionary state of galaxies (e.g., McGaugh & de Blok 1997; Schombert et al. 2001). As an example of the utility of a constant mass-to-light ratio, we adopt $\Upsilon_*^{[3.6]} = 0.5 M_\odot/L_\odot$ in order to plot the gas fraction of galaxies $f_g = M_g/(M_* + M_g)$ as a function of stellar mass in Fig. 7. This is particularly useful for gas rich galaxies that have small but highly uncertain stellar masses. Fig. 7 shows a reasonable distribution of galaxies in the f_g - M_* plane. Simply taking a constant NIR Υ_* appears to give plausible stellar mass estimates, even for objects with formally negative stellar masses according to the BTFR method.

Fig. 7 has a well defined upper envelope. This is easily anticipated from the definition of f_g , which depends on the stellar mass. It is also real: galaxies with large stellar masses and high gas fraction are intrinsically rare. Such objects would be bright in both optical and 21 cm luminosity, and readily detectable in large surveys. Objects with $M_* \approx 10^{11} M_\odot$ have converted most of their original gas into stars by the present epoch. This is consistent with the downsizing of galaxies observed over a range of redshifts, in which massive galaxies typically appear to be the oldest and most evolved at any given epoch. Note, however, that this conclusion would not persist if Υ_* were much smaller, as even the most massive spirals would then be inferred to be rather gas rich.

Galaxies of low stellar mass run the gamut from very gas rich to gas poor. The smallest galaxies discussed here overlap with the classical dwarf Spheroidals of the Local Group in terms of stellar mass. Fig. 7 shows Fornax as an example of such objects. Though the lowest mass galaxies discussed here may have fewer stars than Fornax, they are still larger in terms of baryonic mass




Figure 7. The gas fraction as a function of stellar mass assuming $\Upsilon_*^{[3.6]} = 0.5 \text{ M}_\odot/\text{L}_\odot$. The galaxy sample spans a large range in mass and samples a range of gas fractions at each mass. The absence of galaxies with large stellar masses and large gas fractions is presumably real. Downsizing is evident in that the more massive galaxies have typically consumed a larger fraction of their gas supply. One should however be cautious about the absence of low stellar mass galaxies with low gas fractions: this is presumably a selection effect, as HI must be detected to measure a gas fraction. Low mass but gas poor galaxies like the Fornax dwarf spheroidal lie in the lower left corner of this diagram.

since their gas fractions are high.

The paucity of galaxies in the lower left part of Fig. 7 is almost certainly a selection effect. In order to be included in our sample, a galaxy must have a positive HI detection. Dwarfs like Fornax would not be included even if detected optically. Hence there is every reason to expect that galaxies fill in the space below the upper envelope. Indeed, our current sample does a good job of sampling the available parameter space, in spite of the inevitable cut-off at low masses and low gas fractions.

The mere existence of gas rich disk galaxies with $M_* \approx 10^7 \text{ M}_\odot$ has interesting implications. This is not much larger than a globular cluster. It is widely expected that the early re-ionization of the universe would suppress the accretion of gas into low mass halos (e.g., Ricotti & Gnedin 2005). This is frequently invoked to explain the properties of the dwarf spheroidals of the Local Group, in particular that they have no gas now, nor would have had for most of their history. If re-ionization is the only part of this story (see McGaugh & Wolf 2010), it should hold everywhere: there should be no galaxies as low in mass as the dSph population with lots of gas. Yet there are examples in Figure 7 of objects with stellar masses comparable to Fornax that are replete with gas (see also Tikhonov & Klypin 2009).

6. CONCLUSIONS

We have presented new Spitzer data for disk galaxies spanning ten magnitudes in [3.6] absolute magnitude. These galaxies also span the full range of gas fractions, from $\sim 10\%$ to over 90% . We have used these data to test the self-consistency of stellar population synthesis models from the optical to NIR bands. We have also applied the BTFR to obtain an independent, kinematic estimate of the calibration of stellar mass-to-light ratios.

Our main conclusions can be summarized as follows:

- Many commonly utilized stellar population models

are not self-consistent in the sense that application of the same model to the same galaxy results in different stellar masses depending on whether an optical or NIR luminosity is used. This problem occurs in the sense that the NIR luminosity is overestimated relative to the optical luminosity for a given mass of stars. This problem is present in all *ab initio* models considered, but is particularly severe in models that include TP-AGB stars.

- Self-consistency between optical and NIR observations can be achieved if NIR mass-to-light ratios are approximately constant and do not vary perceptibly with color. The typical value for self-consistency is $0.5 \text{ M}_\odot/\text{L}_\odot$ at $3.6\mu\text{m}$ (equivalent to $0.65 \text{ M}_\odot/\text{L}_\odot$ in the K_s -band).
- The BTFR calibrated by gas dominated galaxies provides an independent constraint on stellar mass. This approach also yields a typical $3.6\mu\text{m}$ mass-to-light ratio of $0.5 \text{ M}_\odot/\text{L}_\odot$.
- The intrinsic scatter about the typical mass-to-light ratio is modest for the most precise data, ~ 0.13 dex in the NIR.
- The IMF is consistent with that of Kroupa (2002).

Both independent methods yield the same typical mass-to-light ratio, giving some hope that this is the correct conversion between light and stellar mass. Within the context of our own study, it appears that the typical value is good to about 20% . We cannot exclude the possibility of systematics at the factor of two level, though these appear unlikely.

Workers wishing to estimate the stellar masses of galaxies would do well to adopt a constant NIR mass-to-light ratio as calibrated here. There appears to be little added value in fitting the complete SED. Indeed, such fits can only be as good as the population model the data are fit to. Given the systematic offsets in the models pointed out here, such SED fits are bound to suffer from systematic errors in stellar mass that will depend on the specific model employed and also on the range of wavelength fit.

This work is based in part on observations made with the Spitzer Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. Support for this work was provided by NASA through an award issued by JPL/Caltech. Other aspects of this work were supported in part by NASA ADAP grant NNX11AF89G and NSF grant AST 0908370. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Begum, A., Chengalur, J. N., Karachentsev, I. D., & Sharina, M. E. 2008, MNRAS, 386, 138
 Bell, E. F., & de Jong, R. S. 2000, MNRAS, 312, 497
 —. 2001, ApJ, 550, 212

- Bell, E. F., McIntosh, D. H., Katz, N., & Weinberg, M. D. 2003, *ApJS*, 149, 289
- Bershady, M. A., Martinsson, T. P. K., Verheijen, M. A. W., Westfall, K. B., Andersen, D. R., & Swaters, R. A. 2011, *ApJ*, 739, L47
- Bershady, M. A., Verheijen, M. A. W., Swaters, R. A., Andersen, D. R., Westfall, K. B., & Martinsson, T. 2010, *ApJ*, 716, 198
- Bottama, R., Pestaña, J. L. G., Rothberg, B., & Sanders, R. H. 2002, *A&A*, 393, 453
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Chabrier, G. 2003, *ApJ*, 586, L133
- Conroy, C., & Gunn, J. E. 2010, *ApJ*, 712, 833
- Dale, D. A., et al. 2005, *ApJ*, 633, 857
- de Blok, W. J. G., McGaugh, S. S., & van der Hulst, J. M. 1996, *MNRAS*, 283, 18
- de Blok, W. J. G., Walter, F., Brinks, E., Trachternach, C., Oh, S.-H., & Kennicutt, R. C. 2008, *AJ*, 136, 2648
- Into, T., & Portinari, L. 2013, *ArXiv e-prints*
- Jarrett, T. H., Chester, T., Cutri, R., Schneider, S., Skrutskie, M., & Huchra, J. P. 2000, *AJ*, 119, 2498
- Kennicutt, Jr., R. C., et al. 2003, *PASP*, 115, 928
- Kriek, M., et al. 2010, *ApJ*, 722, L64
- Kroupa, P. 2002, *Science*, 295, 82
- Leroy, A. K., Walter, F., Brinks, E., Bigiel, F., de Blok, W. J. G., Madore, B., & Thornley, M. D. 2008, *AJ*, 136, 2782
- Maraston, C. 2005, *MNRAS*, 362, 799
- McGaugh, S. S. 2005, *ApJ*, 632, 859
- . 2011, *Physical Review Letters*, 106, 121303
- . 2012, *AJ*, 143, 40
- McGaugh, S. S., & de Blok, W. J. G. 1997, *ApJ*, 481, 689
- . 1998, *ApJ*, 499, 41
- McGaugh, S. S., Schombert, J. M., Bothun, G. D., & de Blok, W. J. G. 2000, *ApJ*, 533, L99
- McGaugh, S. S., & Wolf, J. 2010, *ApJ*, 722, 248
- Melbourne, J., et al. 2012, *ApJ*, 748, 47
- Oh, S., de Blok, W. J. G., Walter, F., Brinks, E., & Kennicutt, R. C. 2008, *AJ*, 136, 2761
- Portinari, L., Sommer-Larsen, J., & Tantaló, R. 2004, *MNRAS*, 347, 691
- Ricotti, M., & Gnedin, N. Y. 2005, *ApJ*, 629, 259
- Schombert, J. 2011, *Astrophysics Source Code Library*, 7011
- Schombert, J., Maciel, T., & McGaugh, S. 2011, *Advances in Astronomy*, 2011
- Schombert, J., & Rakos, K. 2009, *AJ*, 137, 528
- Schombert, J. M., McGaugh, S. S., & Eder, J. A. 2001, *AJ*, 121, 2420
- Stark, D. V., McGaugh, S. S., & Swaters, R. A. 2009, *AJ*, 138, 392
- Swaters, R. A., Sancisi, R., van Albada, T. S., & van der Hulst, J. M. 2009, *A&A*, 493, 871
- Tikhonov, A. V., & Klypin, A. 2009, *MNRAS*, 395, 1915
- Trachternach, C., de Blok, W. J. G., McGaugh, S. S., van der Hulst, J. M., & Dettmar, R. 2009, *A&A*, 505, 577
- Tully, R. B., Rizzi, L., Shaya, E. J., Courtois, H. M., Makarov, D. I., & Jacobs, B. A. 2009, *AJ*, 138, 323
- Verheijen, M. A. W. 2001, *ApJ*, 563, 694
- Walter, F., Brinks, E., de Blok, W. J. G., Bigiel, F., Kennicutt, R. C., Thornley, M. D., & Leroy, A. 2008, *AJ*, 136, 2563
- Zibetti, S., Charlot, S., & Rix, H.-W. 2009, *MNRAS*, 400, 1181
- Zibetti, S., Gallazzi, A., Charlot, S., Pierini, D., & Pasquali, A. 2013, *MNRAS*, 428, 1479